


AD 675499

100 (1)



Carnegie-Mellon University

PITTSBURGH, PENNSYLVANIA 15213

GRADUATE SCHOOL OF INDUSTRIAL ADMINISTRATION

WILLIAM LARIMER MELLON, FOUNDER



Reproduced by the
CLEARINGHOUSE
for Federal Scientific & Technical
Information Springfield Va 22151

//

Management Sciences Research Report No. 127

A NOTE ON LINEAR PROGRAMMING
AND
INTEGER FEASIBILITY

by

Fred Glover^{*}

August, 1968

This report was prepared as part of the activities of the Management Sciences Research Group, Carnegie-Mellon University under Contract NONR 760(24) NR 047-048 with the U. S. Office of Naval Research and the College of Business Administration, Austin, Texas. Reproduction of this paper in whole or in part is permitted for any purpose of the United States Government.

^{*}The University of Texas at Austin, Texas.

Management Sciences Research Group
Graduate School of Industrial Administration
Carnegie-Mellon University
Pittsburgh, Pennsylvania

The zero-one mixed integer programming problem may be written

$$\begin{aligned}
 & \text{Maximize } cx + dy \\
 & \text{subject to} \quad Ax + Dy \leq b \\
 (1) \quad & x \leq e^\dagger \\
 & x, y \geq 0 \text{ and } x \text{ integer}
 \end{aligned}$$

where A is $m \times n$, D is $m \times r$, x is $n \times 1$, y is $r \times 1$, and c , d , b , and e are dimensioned compatibly. Adding vectors u and v of slack variables, the constraints of (1) become

$$\begin{aligned}
 (2) \quad & Ax + Dy + u = b^{\dagger\dagger} \\
 & x + v = e \\
 & x, y, u, v \geq 0 \text{ and } x \text{ integer.}
 \end{aligned}$$

A strategy that is often used for solving (1) is to adjoin additional constraints and variables to partition the feasible region into subsets (e.g., restricting $cx + dy$ to specified intervals), and seek a feasible solution to the additionally constrained problem. We assume that such constraints and variables are already included in (1) and (2), and address ourselves to obtaining feasible solutions to (2).

In a variety of practical situations the constraints of (2) contain imbedded network problems (such as transportation and assignment problems) having the property that every extreme point solution is integer. Moreover, one can readily derive constraints from (2) that impose bounds on nested partial sums of variables (see [3]), and these constraints also have the property that every extreme point is integer. Finally, a set of such constraints can be adjoined to further partition the feasible region of (2).

\dagger e denotes a vector of ones.

$\dagger\dagger$ The results to follow are unchanged if this constraint is more generally $Ax + Dy + uU = b$, provided the augmented matrix (D, U) contains an $m \times m$ non-singular submatrix.

Thus, representing these special constraints by $Px = f$ and $Qx + w = g$, consider the augmented system

$$\begin{aligned}
 (3) \quad & Ax + Dy + u && = b \\
 & x &+ v &= e \\
 & Px &&= f \\
 & Qx &+ w &= g \\
 & x, y, u, v, w \geq 0 \text{ and } x \text{ integer,}
 \end{aligned}$$

where P , Q , f and g are integer matrices, P is $p \times n$, Q is $q \times n$, and $\begin{pmatrix} D \\ Q \end{pmatrix}$ has the unimodular property; i.e., every square submatrix of $\begin{pmatrix} D \\ Q \end{pmatrix}$ has determinant 0, 1, or -1. (P or Q may also be null.) We also stipulate $p \leq n$ and every $p \times p$ submatrix of P is nonsingular.

Theorem. If there is a feasible solution to (3) with $x = x'$ and x' integer, then there is a basic feasible solution to (3) with $x = x'$ and m of the components of (y, u) basic. Moreover, every basic feasible solution to (3) with m of the components of (y, u) basic assigns integer values to the components of x , v , and w .

The chief significance of this theorem is that it permits one to elect a strategy for solving (2) that focuses on finding a solution to (3) with m of the components of (y, u) basic.

The first application of such a strategy occurs in the pure zero-one linear programming method of Cabot and Hurter [1], whose ideas motivate this note. Specifically, the Cabot and Hurter method results (for D and d null) by adjoining the constraint $ex = N$ to (2) and replacing b by $b + \epsilon e$, where $0 < \epsilon < 1$ and A and b are assumed integer. Beginning with N at an upper bound for ex , Cabot and Hurter prescribe finding a basic feasible solution to this particular version of (3) with all components of u basic. (No procedure is given for

accomplishing this, however.) If an acceptable solution is found, the method stops (or is applied to a new system (2)). Otherwise, N is decremented and the process repeats.

The theorem implies that $ex = N$ can be dispensed with in the Cabot and Hurter approach, making it unnecessary to reapply the process for different values of N . Moreover, a variety of other side conditions and supplementary constraints can be accommodated by the theorem, some of the more important of which have already been indicated.

A method for exploiting the theorem for the pure integer programming problem is given below. We first establish the validity of the theorem with the following three lemmas.

Lemma 1. Every square submatrix of

$$H = \begin{pmatrix} I & I & 0 \\ P & 0 & 0 \\ Q & 0 & I \end{pmatrix} \text{ has determinant } 0, 1, \text{ or } -1$$

(where the I matrices of the top row are $n \times n$).

Proof: First, it is assumed that $\begin{pmatrix} P \\ Q \end{pmatrix}$ has the unimodular property. Using induction on q , and expanding the determinants of the appropriate submatrices of H by minors, it is easy to see that $\begin{pmatrix} P & 0 \\ Q & I \end{pmatrix}$ has the unimodular property. The rest of the proof follows the same argument, using induction on n .

Lemma 2. Consider the system

$$Mt + Rx = \alpha$$

$$Hz = \beta \quad \text{and } t, z \geq 0$$

where M is $m \times l$, R is $m \times s$, and H is $h \times s$, the vectors α and β dimensioned compatibly. If β is integer and every square submatrix of H has determinant 0, 1 or -1, then z is integer in every basic solution to (4) with m of the components of t basic.

Proof: The basic solution must have the form

$t = M_1^{-1} (\alpha - R_1 H_1^{-1} \beta)$ and $z = H_1^{-1} \beta$, where M_1 is an $m \times m$ submatrix of M , R_1 is an $m \times h$ submatrix of R , and H_1 is an $h \times h$ submatrix of H . The latter assures z is integer (see Hoffman and Kruskal [5]).

Lemmas 1 and 2 collectively imply the latter part of the theorem. The first part of the theorem is implied by the following stronger statement.

Lemma 3. If there is a feasible solution to (3) with $x = x'$ and x' integer then *and x' integer* there is a basic feasible solution to (3) in which

- (i) every component of w is basic,
- (ii) x_j is basic if $x'_j = 1$ and v_j is basic if $x'_j = 0$
- (iii) any p of the remaining components of (x, v) are basic,
- (iv) m of the components of (y, u) are basic.

Proof: Conditions (i), (ii), (iii) give $q + n + p$ variables. We show that the submatrix composed of the associated columns of H of Lemma 1 (call it H_1) is nonsingular. First, all columns of Q in H , may be reduced to 0 by subtracting from them appropriate multiples of the columns of $I_{q \times q}$ (associated with w).

Next, there must be exactly p indices j such that x_j and v_j are chosen to be basic. Subtracting each of these v_j columns from its associated x_j column, and rearranging rows and columns, transforms H_1 into $\begin{pmatrix} P_1 & R \\ 0 & I \end{pmatrix}$, where P_1 is a $()$ $p \times p$ submatrix of P , R consists of 0 columns and columns of P , and I is the $(n+q) \times (n+q)$ identity matrix. The nonsingularity of P_1 assures H_1 is nonsingular, and hence a basis for the subsystem of (3) with $Ax + Dy + u = b$ removed. Since H_1 contains a column for each positive component of x' , t' , and w' (where $t' = e - x'$, $w' = g - Qx'$), the basic solution to the subsystem must yield $x = x'$. Finally, (iv) is established in conjunction with (i), (ii), and (iii) from the assumed existence of a feasible, and hence a basic feasible, solution to $Dy + u = b - Ax'$.

An Integer Programming Method

We give an integer programming method for the pure zero-one problem (with D and d null) that pursues the objective of making the m components of u basic in (3). Assume A and b are integer, and replace A by $2A$ and b by $2b + e$. (This replacement clearly does not change the set of integer x satisfying $Ax \leq b$, and implies $u \geq e$ and integer for all non-negative u , x satisfying $u + Ax = b$ and x integer.)

1. Solve the linear program

$$\text{Maximize } x_0 = au$$

subject to (3) (disregarding the integer restriction on x),

where $a > 0$ and integer (e.g., let $a = e$).

Represent the current tableau for the simplex method in the form

$$\begin{aligned} \text{Maximize } x_0 &= a_{00} + \sum_{j=1}^l a_{0j} (-t_j) \\ z_i &= a_{i0} + \sum_{j=1}^l a_{ij} (-t_j) \quad i = 1, \dots, p \end{aligned}$$

where the z_i are the current basic variables and the t_j the current nonbasic variables. (The a_{ij} coefficients of the current tableau are not to be confused with the components of the A matrix.)

Upon obtaining an optimal tableau ($a_{i0} \geq 0$ and $a_{0j} \geq 0$ for $i, j \geq 1$), go to Step 2.

2. If a_{i0} is integer for all i , the basic solution $z_i = a_{i0}$ and $t_j = 0$ (all i, j) gives a feasible integer solution for (2) by identifying the variables x_j from among those currently designated z_i and t_j . Otherwise, if some a_{i0} is noninteger, adjoin a cut [4, 2] and re-optimize with the dual simplex method.

3. If some a_{10} is still noninteger, let t_s denote the current nonbasic variable that was the slack variable for the cut adjoined (most recently) in Step 2. Replace a_{os} with $a_{os} - K < 0$, where K is an integer (e.g., the least integer $< a_{os}$). Then re-optimize with the primal simplex method and return to 2.

The purpose of Step 3 is to exploit the fact that the cut slack qualifies to be one of the " u_1 " of the theorem. Thus, it assigns the slack a weight in the objective function designed to drive it basic, thereby possibly modifying, but not discarding, the weights assigned to the other u_1 .

Finite convergence is guaranteed if one uses the choice rules of Gomory [4] in Step 2 and bypasses Step 3 after a fixed number of iterations. The main point, of course, is that the method gives a way to pursue integer feasibility by exploiting the theorem.

Acknowledgment: I am indebted to M. Raghavachari of Carnegie-Mellon University for his very constructive criticism of this note.

References

- [1] Cabot, A. V., and Arthur P. Hurter, Jr., "An algorithm for zero-one integer programming,"
- [2] Glover, Fred, "Generalized cuts in diophantine programming," Management Science, Vol. 13, No. 3, Nov, 1966.
- [3] _____, "Flows in arborescences," Management Sciences Research Report No. 102, Carnegie-Mellon University, July, 1967.
- [4] Gomory, Ralph, "An algorithm for integer solutions to linear programs," in R. L. Graves and Ph. Wolfe (eds.), Recent Advances in Mathematical Programming, McGraw Hill, 1963.
- [5] Hoffman, A. J., and J. B. Kruskal, "Integral Boundary Points of Convex Polyhedra," in Kuhn and Tucker (eds.), Annals Study 38, Linear Inequalities and Related Systems, Princeton University Press, 1956.

Unclassified

Security Classification

DOCUMENT CONTROL DATA - R & D		
(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)		
1. ORIGINATING ACTIVITY (Corporate author) Graduate School of Industrial Administration Carnegie-Mellon University		2a. REPORT SECURITY CLASSIFICATION Unclassified
		2b. GROUP Not applicable
3. REPORT TITLE A Note on Linear Programming and Integer Feasibility		
4. DESCRIPTIVE NOTES (Type of report and, inclusive dates) Technical Report, August, 1968		
5. AUTHOR(S) (First name, middle initial, last name) Fred Glover		
6. REPORT DATE August, 1968	7a. TOTAL NO. OF PAGES 9	7b. NO. OF REFS 5
8a. CONTRACT OR GRANT NO. NONR 760(24)	8b. ORIGINATOR'S REPORT NUMBER(S) Management Sciences Research Report No. 127	
b. PROJECT NO. NR 047-048		
c.	9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report) Not applicable	
d.		
10. DISTRIBUTION STATEMENT Distribution of this document is unlimited		
11. SUPPLEMENTARY NOTES Also under activities of the University of Texas at Austin		12. SPONSORING MILITARY ACTIVITY Logistics and Mathematical Statistics Br. Office of Naval Research, Washington, D.C.
13. ABSTRACT A theorem is given that provides new strategies for solving integer programming problems, based on finding certain types of basic solutions to linear programs. The theorem is motivated by and extends ideas of Cabot and Hurter [1]. An integer programming method based on the theorem is outlined.		

DD FORM 1473

1 NOV 68
S/N 0101-807-6811

(PAGE 1)

Unclassified

Security Classification

A-31408

Security Classification

KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
Integer Programming						
Linear Programming						
Zero-One Integer Programming						